

---

# What Are Radicals In Math

---

*What Are Radicals In Math*

Downloaded from  
[db.whlarchitecture.com](http://db.whlarchitecture.com) by Elaine &  
Callahan

---

## INTRODUCTION: UNLOCKING THE LANGUAGE OF ROOTS AND RADICALS

---

Mathematics often feels like a foreign language, and one of its most intriguing characters is the radical symbol:  $\sqrt{\phantom{x}}$ . You have likely seen it in geometry when calculating the diagonal of a square, or in algebra when solving for an unknown variable. But what exactly is a radical? At its core, a radical represents the "inverse" operation of exponentiation. If squaring a number means multiplying it by itself, then taking a radical—specifically a square root—asks the question: "What number, when multiplied by itself, gives me this value?"

However, radicals are much more than just square roots. They include cube roots, fourth roots, and beyond. Understanding radicals is not just an academic exercise; it is a foundational skill that unlocks advanced topics in algebra, calculus, physics, engineering, and even computer science. Whether you are a student struggling with homework, a parent trying to help, or a professional brushing up on fundamentals, this guide will walk you through everything you need to know about radicals in a

clear and intuitive way.

---

## WHAT IS A RADICAL? BREAKING DOWN THE NOTATION

---

A radical expression is defined by three key components: the **radical symbol** ( $\sqrt{\phantom{x}}$ ), the **radicand** (the number or expression inside the radical), and the **index** (the small number written just outside and above the radical sign). The index tells you which root you are taking. If no index is written, it is assumed to be 2, which means the square root.

For example, in the expression  $\sqrt[3]{27}$ , the index is 3, the radicand is 27, and the radical symbol indicates that we are looking for the cube root of 27. The answer is 3 because  $3 \times 3 \times 3 = 27$ . In the expression  $\sqrt{36}$ , the index is 2 (implied), the radicand is 36, and the answer is 6 because  $6 \times 6 = 36$ .

Understanding this notation is the first step. But radicals are not just about numbers; they can also contain variables. For instance,  $\sqrt{x^2 + 4}$  asks for the square root of the entire expression. This is where radicals become powerful tools for solving equations and modeling real-world scenarios.

---

## EXPONENTS

To truly grasp what radicals are, you must understand their relationship with exponents. A radical is simply another way of writing a fractional exponent. This connection is not just a mathematical curiosity; it is the key to simplifying complex expressions and solving equations efficiently.

The general rule is: the  $n$ th root of a number is equivalent to raising that number to the power of  $1/n$ . In symbolic form,  $\sqrt[n]{a} = a^{(1/n)}$ . For example, the square root of  $x$  is  $x^{(1/2)}$ , and the cube root of  $y$  is  $y^{(1/3)}$ . This equivalence allows you to switch between radical form and exponent form, often making calculations easier.

Consider the expression  $\sqrt{(x^6)}$ . Instead of working with the radical, you can rewrite it as  $(x^6)^{(1/2)}$ . Using the power rule of exponents, you multiply the exponents:  $6 \times 1/2 = 3$ , so the expression simplifies to  $x^3$ , provided  $x$  is non-negative. This technique is especially useful when dealing with higher-level algebra and calculus.

Conversely, any expression with a fractional exponent can be converted into a radical. For instance,  $a^{(2/3)}$  is the same as  $(\sqrt[3]{a})^2$  or  $\sqrt[3]{(a^2)}$ . Both forms are valid, and choosing the right one depends on the context of the problem. Mastering this interchange is essential for advancing in mathematics.

## TYPES OF RADICALS: BEYOND THE SQUARE ROOT

While square roots are the most common type of radical, there is

a whole family of radicals to be explored. Understanding the different types helps you see the bigger picture and avoid confusion.

## Square Roots

The square root of a number  $a$  is the value that, when multiplied by itself, equals  $a$ . The symbol is  $\sqrt{a}$ . For example,  $\sqrt{16} = 4$  because  $4 \times 4 = 16$ . Square roots are fundamental in geometry (think of the Pythagorean theorem), physics (calculating velocity or energy), and everyday calculations.

## Cube Roots

The cube root of a number  $a$  is the value that, when multiplied by itself three times, equals  $a$ . The symbol is  $\sqrt[3]{a}$ . For example,  $\sqrt[3]{64} = 4$  because  $4 \times 4 \times 4 = 64$ . Cube roots are particularly useful in volume calculations and in solving cubic equations. Unlike square roots, cube roots can be taken of negative numbers.  $\sqrt[3]{(-8)} = -2$  because  $(-2) \times (-2) \times (-2) = -8$ .

## Fourth Roots and Beyond

Fourth roots, fifth roots, and  $n$ th roots follow the same pattern. The fourth root of a number  $a$  is the value that, when multiplied by itself four times, equals  $a$ . The symbol is  $\sqrt[4]{a}$ . For example,  $\sqrt[4]{81} = 3$  because  $3 \times 3 \times 3 \times 3 = 81$ . For even roots (square, fourth, sixth, etc.), the radicand must be non-negative if you are working with real numbers. For odd roots (cube, fifth, seventh,

etc.), the radicand can be any real number.

---

### SIMPLIFYING RADICALS: THE ART OF BREAKING DOWN

---

Simplifying radicals is one of the most important skills to develop. A radical is considered "simplified" when the radicand has no perfect square factors (for square roots), no perfect cube factors (for cube roots), and so on. The goal is to make the expression as clean and manageable as possible.

## Simplifying Square Roots

To simplify  $\sqrt{72}$ , you look for the largest perfect square that divides 72. The perfect squares are 4, 9, 16, 25, 36, etc. Since 36 divides 72 ( $36 \times 2 = 72$ ), you can rewrite  $\sqrt{72}$  as  $\sqrt{(36 \times 2)}$ . Then, using the property  $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$ , you get  $\sqrt{36} \times \sqrt{2} = 6\sqrt{2}$ . So the simplified form of  $\sqrt{72}$  is  $6\sqrt{2}$ .

Here is a step-by-step process for simplifying any square root:

1. **Factor the radicand** into its prime factors or find the largest perfect square factor.
2. **Rewrite the radical** as the product of two radicals, one with the perfect square.
3. **Take the square root** of the perfect square factor.
4. **Multiply** the result by the remaining radical.

## Simplifying Cube Roots and Higher

The same principle applies to cube roots, but you look for perfect cube factors (1, 8, 27, 64, 125, etc.). For example,  $\sqrt[3]{54} = \sqrt[3]{(27 \times 2)} = \sqrt[3]{27} \times \sqrt[3]{2} = 3 \times \sqrt[3]{2} = 3\sqrt[3]{2}$ . For fourth roots, look for perfect fourth powers like 16, 81, 256, and so on.

---

### OPERATIONS WITH RADICALS

---

Just like numbers and variables, radicals can be added, subtracted, multiplied, and divided. However, there are specific rules that must be followed to ensure accuracy.

## Adding and Subtracting Radicals

You can only add or subtract radicals that have the **same index** and the **same radicand**. These are called "like radicals," analogous to like terms in algebra. For example,  $3\sqrt{5} + 2\sqrt{5} = 5\sqrt{5}$ . However,  $\sqrt{3} + \sqrt{2}$  cannot be simplified further because the radicands are different. If the radicals are not alike, you must first simplify them and then check again for common radicands.

# Multiplying Radicals

Multiplying radicals is straightforward as long as the indices are the same. The rule is:  $\sqrt[n]{a} \times \sqrt[n]{b} = \sqrt[n]{ab}$ . For example,  $\sqrt{3} \times \sqrt{12} = \sqrt{36} = 6$ . If the indices are different, you must first convert the radicals to exponential form, find a common denominator for the exponents, and then multiply.

# Dividing Radicals and Rationalizing the Denominator

Dividing radicals follows a similar rule:  $\sqrt[n]{a} \div \sqrt[n]{b} = \sqrt[n]{a/b}$ , provided  $b \neq 0$ . For example,  $\sqrt{18} \div \sqrt{2} = \sqrt{9} = 3$ .

A special case arises when a radical appears in the denominator of a fraction. It is considered standard mathematical practice to **rationalize the denominator**, which means rewriting the fraction so that the denominator no longer contains a radical. For a simple case like  $1/\sqrt{2}$ , you multiply the numerator and denominator by  $\sqrt{2}$  to get  $\sqrt{2}/2$ . For more complex denominators, such as those containing binomials like  $1/(\sqrt{3} + 1)$ , you multiply by the conjugate  $(\sqrt{3} - 1)$  to eliminate the radical.

## RADICALS AND RATIONAL EXPONENTS: A POWERFUL

## PARTNERSHIP

As mentioned earlier, radicals and rational exponents are two sides of the same coin. Understanding this relationship allows you to simplify expressions that would otherwise be cumbersome. For instance, consider the expression  $\sqrt[5]{x^{10}}$ . Using exponent notation, this is  $(x^{10})^{(1/5)} = x^{(10/5)} = x^2$ . This is much faster than trying to factor the radical.

Rational exponents also allow you to combine operations. For example,  $(\sqrt[3]{x})^4$  can be written as  $x^{(4/3)}$ . This flexibility is invaluable when solving equations or working with functions. When you encounter a variable raised to a rational exponent, you can often rewrite it as a radical to help visualize the root being taken.

## COMMON MISTAKES AND HOW TO AVOID THEM

Every student encounters pitfalls when learning about radicals. Being aware of these common errors can save you time and frustration.

- **Mistake 1: Treating  $\sqrt{a + b}$  as  $\sqrt{a} + \sqrt{b}$ .** This is false. For example,  $\sqrt{(9 + 16)} = \sqrt{25} = 5$ , but  $\sqrt{9} + \sqrt{16} = 3 + 4 = 7$ . Addition does not distribute over the radical.
- **Mistake 2: Forgetting the index.** Students often treat every radical as a square root, even when the index is 3 or higher. Always check the index carefully.
- **Mistake 3: Ignoring absolute values with even roots.** When taking an even root (like a square root) of a variable squared, the result is the absolute value of the variable. For

example,  $\sqrt{x^2} = |x|$ , not just  $x$ , because the square root always returns a non-negative result.

- **Mistake 4: Not simplifying fully.** Always check if the radicand has any perfect power factors (squares, cubes, etc.) before declaring your final answer.

---

### REAL-WORLD APPLICATIONS OF RADICALS

---

Radicals are not just theoretical concepts; they appear in many practical fields. Understanding them opens doors to solving real problems.

## Geometry and Physics

The Pythagorean theorem, which states that in a right triangle,  $a^2 + b^2 = c^2$ , often requires taking a square root to find the

hypotenuse  $c$ . This is used in construction, navigation, and computer graphics. In physics, the formula for the period of a pendulum involves a square root, as does the formula for the velocity of a falling object.

## Finance and Economics

Compound interest formulas and the calculation of root mean square (RMS) are applications of radicals. The RMS value is used in electrical engineering to measure alternating current (AC) voltage, and it involves taking the square root of the mean of squares.

## Medicine and Biology

Radicals